Chicago, IllinoisžUSA New World Blackbirds (Icteridae) of Greater Grant Park

Michelle Buziak Field Museum

Produced and designed by Michelle Buziak. Acknowledgments: Ben Marks and the Field Museum Bird Collections staff; Greater Grant Park Bird List contributors Doug Stotz, Dave Willard, and Josh Engel. University of California Scientific Illustration program, Ann Caudle, Jenny Keller, Amadeo Bachar, Andrea Dingeldein; and Justine Lee Hirten. iNaturalist and their contributors. Illustration Indicates sex: M = Male, F= Female.



CC0. This work is dedicated to the public domain.

[fieldguides.fieldmuseum.org]

[1605]

version 1

6/2023



New World Blackbirds or Icteridae are found in a diverse range of habitats, from open woodlands to marshes to grasslands. The species found in the Chicagoland area are mostly migratory where all species can be found from April to August. The Brown-headed Cowbird, Red-winged Blackbird, Eastern Meadowlark, Brewer's Blackbird and the Common Grackle are found almost year-round except for the coldest months (January-February). The Rusty Blackbird migrates north in the late winter/early spring and returns south in the fall.1


The tree-covered plazas of Millenium Park and the museum campus are great places to look for these birds. Grassland-loving species, such as the Bobolink, can be found in the summer months at Northerly Island Preserve.2

1. Brock, Kenneth J. "Appendix A:." In Birds of the Indiana Dunes, 232-43. N.p., 1997.

Yellow-headed Blackbird

Tordo Cabeciamarillo Xanthocephalus xanthocephalus

2. iNaturalist. Available from https://www.inaturalist.org. Accessed [February 25,2022].



10

Rusty Blackbird

Zanate Canadiense

Euphagus carolinus